

Dedication. Compassion. Experience.

Sean E. Rockett, M.D. Agam A. Shah, M.D. Arthur F. Christiano, M.D. Mark Rose, D.P.M. Ethan M. Healy, M.D. Daniel B. Osuch, M.D. Symeon V. Zannikos, M.D. Mark A. Finno, M.D. Kevin P. Sullivan, M.D. Jason C. Saillant, M.D. Michael L. Sganga, D.P.M. Daren Bergman, D.P.M. David M. Ramsden, M.D.

Nury Flynn, PA-C, ATC Asimina Markopoulos, PA-C Karl Haywood, PA-C Ryan C. Gulla, PA-C Brian Sheehy, PA-C Matthew Richardson, PA-C Hannah Moomey, PA-C Brianna Leung, PA-C

Post Operative Foot Surgical Instruction

1. Activity should be only as recommended by your surgeon as

	follows:
2.	Keep the operative foot elevated on two pillows whenever possible.
3.	Apply ice just above the operative site (usually to the ankle), 30 minutes out of every one
	hour for the first 48 hours after surgery.
4.	Avoid prolonged periods of standing or sitting with legs hanging down.
5.	Keep bandages CLEAN and DRY. Do not remove the bandage.
6.	Never allow the bandage to get wet.
7.	Do not bump or injure your foot.
8.	Keep pets away from the foot.
9.	Take all medications exactly as prescribed.
10.	When you arrive home call the office for an appointment onThe first dressing
	change, foot check and post operative x-rays will occur at this appointment.
11.	You have just had a surgical procedure. A limited amount of swelling is expected. In
	some cases, the skin may take on a bruised appearance. This should not be cause for
	alarm.
12.	If you should notice any increase in temperature, pain, swelling,
	Redness or bleeding through your dressing you should contact your surgeon
	immediately.
	YOUR cooperation determines the successful outcome of your surgery.
	Additional instructions:







2000 Washington Street

Newton