

Dedication. Compassion. Experience.

Sean E. Rockett, M.D. Agam A. Shah, M.D. Arthur F. Christiano, M.D. Mark Rose, D.P.M. Ethan M. Healy, M.D. Daniel B. Osuch, M.D. Symeon V. Zannikos, M.D. Mark A. Finno, M.D. Kevin P. Sullivan, M.D. Jason C. Saillant, M.D. Michael L. Sganga, D.P.M. Daren Bergman, D.P.M. David M. Ramsden, M.D. Nury Flynn, PA-C, ATC Asimina Markopoulos, PA-C Karl Haywood, PA-C Ryan C. Gulla, PA-C Brian Sheehy, PA-C Matthew Richardson, PA-C Hannah Moomey, PA-C Brianna Leung, PA-C

Post-Operative Care Distal Biceps Tendon Repair

Dressing/Wound Care

- Do NOT get the dressing or splint wet. Cover with a plastic bag or plastic wrap for showers.
- You may notice some drainage around the wound, this is normal.
 - o If this persists more than five days, please call the office as a precaution. This can still be normal.
- You may remove the sling only to shower- splint and dressing stays on and in placekeep your arm at your body's side for comfort.
- Some swelling or bruising around the arm, elbow or hand is expected and can last up to several weeks.
- Use ice over your arm for the first 24-36 hours after surgery. You may place ice over the dressing for as long as you feel comfortable (recommend at least 20 min./hour).
 Do not place ice or ice pack in direct contact with skin.
- Low-grade temperatures up to 101 are common after surgery, please call if temperature rises above 101.4.

Ambulation and Movement

- Immediately after surgery you will be placed in a shoulder sling with a splint and bulky dressing.
- Do not remove the dressing or splint- leave intact- this will be removed at your first post-op visit in the office.
- Elevate your hand to sky when sitting, when walking use the sling.
- Perform gentle finger and hand range-of-motion several times per day to prevent stiffness.











Hopkinton



Sean E. Rockett, M.D. Agam A. Shah, M.D. Arthur F. Christiano, M.D. Mark Rose, D.P.M. Ethan M. Healy, M.D. Daniel B. Osuch, M.D. Symeon V. Zannikos, M.D. Mark A. Finno, M.D. Kevin P. Sullivan, M.D. Jason C. Saillant, M.D. Michael L. Sganga, D.P.M. Daren Bergman, D.P.M. David M. Ramsden, M.D. Nury Flynn, PA-C, ATC Asimina Markopoulos, PA-C Karl Haywood, PA-C Ryan C. Gulla, PA-C Brian Sheehy, PA-C Matthew Richardson, PA-C Hannah Moomey, PA-C Brianna Leung, PA-C

Post-Operative Care Distal Biceps Tendon Repair (continued)

Medication Take as prescribed

- You will have received a prescription for pain medication.
 - o Do not operate machinery or drive while taking this medication.
 - This medication may cause nausea or vomiting.
- You may have been given a prescription for an anti-nausea medication.
 - o Take as directed for any nausea associated with the pain medication.
- For added pain control, begin taking Tylenol 500mg tablets, 1-2 tablets every 6 hours as needed.
- Any severe itching, hives or difficulty breathing, please call immediately.
 - Any severe itching, hives or difficulty breathing, please call immediately.

Physical Therapy

- On your first visit after surgery you will be given a prescription for physical therapy.
- This will be one of the important steps leading to your recovery.

Return Visit

- If you do not already have a scheduled post-operative visit please call for an appointment on the next business day.
- The first post-operative visit is usually 7-10 days after date of surgery unless instructed otherwise.
- If you have any problems, do not wait for an appointment, please call the office.







313 Speen Street

Natick



77 West Main Street

Hopkinton